

# Aspirin-triggered 15-epi-lipoxin A<sub>4</sub> regulates neutrophil-platelet aggregation and attenuates acute lung injury in mice

Guadalupe Ortiz-Muñoz<sup>1</sup>, Beñat Mallavia<sup>1</sup>, Adriaan Bins<sup>2</sup>, Mark Headley<sup>3</sup>, Matthew F. Krummel<sup>3</sup>, Mark R. Looney<sup>1,4</sup>

<sup>1</sup>Departments of Medicine, <sup>3</sup>Pathology, and <sup>4</sup>Laboratory Medicine, University of California, San Francisco, San Francisco, CA, USA. <sup>2</sup>Department of Immunology, Netherlands Cancer Institute, Amsterdam, Netherlands.

**Corresponding Author:** Mark R. Looney, M.D.  
513 Parnassus Avenue, HSE 1355A  
San Francisco, CA 94143-0130  
[mark.looney@ucsf.edu](mailto:mark.looney@ucsf.edu)  
Tel: (415) 476-9563  
Fax: (415) 502-2605

**Abbreviated title:** Neutrophil-platelet aggregates and lung injury

**Key words:** neutrophils, platelets, lipoxins, aspirin, acute lung injury, transfusion-related acute lung injury

**Scientific Category:** Platelets and Thrombopoiesis

## **Key Points**

- Neutrophil-platelet aggregates are dynamically formed in the lung in response to injury and are regulated by aspirin-triggered lipoxin.
- The therapeutic effect of aspirin in acute lung injury is in large part mediated by the production of pro-resolving lipid mediators.

## Abstract

Evidence is emerging that platelets are major contributors to innate immune responses in conditions such as acute lung injury (ALI). Platelets form heterotypic aggregates with neutrophils, and we hypothesized that lipoxin mediators regulate formation of neutrophil-platelet aggregates (NPA), and that NPA significantly contribute to ALI. LPS-induced lung injury was accompanied by platelet sequestration, activation, and intra-alveolar accumulation, and NPA formation within both blood and alveolar compartments. Using lung intravital microscopy, we observed the dynamic formation of NPA during physiologic conditions, which sharply increased with ALI. Aspirin treatment significantly reduced lung platelet sequestration and activation, NPA formation, and lung injury. Aspirin treatment increased levels of aspirin-triggered lipoxin (ATL, 15-epi-LXA<sub>4</sub>) and blocking the lipoxin A<sub>4</sub> receptor (ALX) with a peptide antagonist (Boc2) or using ALX knockouts (*Fpr2/3<sup>-/-</sup>*) reversed this protection. LPS increased NPA formation *in vitro*, which was reduced by ATL, and engagement of ALX by ATL on both neutrophils and platelets was necessary to prevent aggregation. In a model of transfusion-related acute lung injury (TRALI), Boc2 also reversed aspirin protection, and treatment with ATL in both LPS and TRALI models protected from ALI. We conclude that ATL regulates neutrophil-platelet aggregation, and that platelet-neutrophil interactions are a therapeutic target in lung injury.

## Introduction

The acute respiratory distress syndrome (ARDS) is a common and often fatal syndrome characterized by neutrophilic inflammation. In ARDS, and experimental models of acute lung injury (ALI), an exuberant inflammatory response with release of proteases and other toxic mediators from activated leukocytes leads to tissue damage, increased permeability of the alveolar-capillary barrier and the formation of protein-rich lung edema.<sup>1</sup>

Several lines of evidence indicate that activated platelets act as important effectors in host defense, influencing pulmonary neutrophil recruitment and contributing to the development of ALI.<sup>2-4</sup> Platelets become activated within the blood circulation at sites of inflammation and can adhere to other platelets, to leukocytes, forming neutrophil-platelet aggregates (NPA), or monocyte-platelet aggregates (MPA), and to exposed endothelium.<sup>5</sup> The presence of circulating leukocyte-platelet aggregates (LPA) is a sensitive indicator of platelet activation and has been observed in patients with inflammatory conditions including sepsis, coronary diseases, rheumatoid arthritis, inflammatory bowel disease, cystic fibrosis, and ARDS.<sup>6-10</sup>

Activated platelets synthesize thromboxane A<sub>2</sub> from arachidonic acid via a sequential reaction catalyzed by cyclooxygenase-2 (COX-2). Aspirin (acetylsalicylic acid, ASA) acetylates COX-2, and this conformational change leads to inhibition of prostanoid synthesis.<sup>11</sup> Acetylation of COX-2 switches catalytic activity to convert arachidonic acid to 15R-hydroxyeicosatetraenoic acid (15R-HETE), which can be subsequently converted to 15(R)-epi-LXA<sub>4</sub>, also known as aspirin-triggered lipoxin (ATL).<sup>12</sup> Lipoxins are endogenous lipid mediators generated during inflammation that can block inflammatory cell recruitment, inhibit cytokine release, and decrease vascular permeability, which collectively are anti-inflammatory properties.<sup>13,14</sup> Lipoxins also stimulate epithelial reconstitution, facilitate apoptosis, and promote macrophage efferocytosis,

which collectively are pro-resolution properties.<sup>13,14</sup> Production of lipoxins requires transcellular biosynthesis, with leukocytes, platelets, and tissue resident cells having important roles.<sup>12,15</sup>

Aspirin-triggered lipoxin (ATL) shares the potent anti-inflammatory actions of lipoxins but is more resistant to metabolic inactivation.<sup>16</sup> The anti-inflammatory, pro-resolving properties of both LXA<sub>4</sub> and 15(R)-epi-LXA<sub>4</sub> are mediated through the formyl peptide receptor type 2 (FPR2, termed Fpr2/3 in the mouse), also called the lipoxin A<sub>4</sub> receptor (ALX),<sup>17</sup> which is expressed in a variety of cell types including endothelial cells and myeloid lineages.<sup>17,18</sup> In addition to native lipoxins and ATL, ALX recognizes a diverse array of bioactive mediators including annexin A1, resolvin D1, and SAA.<sup>17,19</sup> Recent work has elucidated the molecular mechanisms behind the dual actions of ALX (pro- vs. anti-inflammatory responses), which is determined by unique receptor dimerization patterns induced by specific ligands.<sup>20</sup>

Here, we demonstrate the role of ATL in regulating leukocyte-platelet aggregation, and consequently lung injury, in both direct and indirect experimental models of ALI: Intratracheal instillation of lipopolysaccharide (LPS) and a two-event model of transfusion-related acute lung injury (TRALI).<sup>21</sup> We present evidence that platelets have a central role in producing pulmonary inflammation and injury based on critical interactions with neutrophils that are regulated by ATL.

## Methods

*Reagents.* The following reagents were used: aspirin (acetylsalicylic acid); LPS from *Escherichia coli* O55:B5 (Sigma-Aldrich); Boc2 (Genway); 15-epi-LXA<sub>4</sub> (EMD Biosciences)

*Intratracheal LPS model.* Experimental procedures were performed in 8-12-week-old male BALB/c wild-type mice (Charles River) housed under pathogen-free conditions in the Animal Barrier Facility at UCSF. Mice were pre-treated (i.p.) with 0.1 mg/g of aspirin or DMSO-vehicle, 24h and 2h before ALI induction. Mice were challenged with intratracheal PBS (controls) or LPS (5 µg/g) and analysis was performed 4- to 48h later. In selected experiments, mice received 10 µg/kg i.p. of Boc2 30min before ASA injections and 24h after LPS instillation, or 100-5000 ng/mouse of 15-epi-LXA<sub>4</sub> 30min before and 24h after LPS challenge. There is no appropriate peptide control for Boc2. *Fpr2/3*<sup>-/-</sup> mice were obtained from Rod Flower (St. Barts and the London School of Medicine and Dentistry).<sup>19</sup>

Bronchoalveolar lavage (BAL) was obtained at 48h by placing a catheter into the trachea, through which 1 ml of cold PBS was flushed back x 3. Protein concentration in the cell-free BAL was determined using a BCA protein assay kit (Thermo Scientific). BAL leukocytes were counted using a Coulter counter (Beckman Coulter), and differential was determined by cytopspin preparation (Cytospin 3, Thermo Electron Corp.) and Diff-Quick staining. BAL hemoglobin was measured using LowHb microcuvettes (HemoCue).

*TRALI model.* A two-event TRALI model was used, as previously described.<sup>21</sup> Briefly, after being primed with LPS (0.1 mg/kg, i.p.) for 24 hours, mice were challenged with MHC I mAb (H2K<sup>d</sup>; IgG<sub>2a</sub>, κ; 1.0 mg/kg) injected into the jugular vein, and euthanized after 2 hours. To quantify pulmonary edema formation, bloodless, extravascular lung water was measured.<sup>21,22</sup> We also measured lung vascular permeability to protein by instilling mice with i.v. <sup>125</sup>I-labeled

albumin. The radioactivity in the blood and the bloodless lung was measured with a gamma counter (Packard 5000 Series), and the ratio was used to calculate the lung extravascular plasma equivalents (EVPE).<sup>21,22</sup> ASA (0.1 mg/g), or DMSO control, was delivered i.p. 30 minutes prior to LPS priming, and again 2h prior to H2K<sup>d</sup> mAb challenge. Boc2 (10 µg/kg), or vehicle (PBS), was administered i.p. 30 min prior to the ASA injections. 15-epi-LXA<sub>4</sub>, or vehicle (ethanol), was administered via tail vein 30 min prior to H2K<sup>d</sup> mAb injections.

*Immunohistochemistry.* For histological analysis, 7µm of OCT-embedded frozen lung sections (without lung perfusion) were fixed in cold acetone and blocked with PBS containing 8% rabbit serum and 4% BSA. Sections were incubated with CD41 antibody (clone MWReg30, BD Biosciences), followed by peroxidase-conjugated secondary Abs, color development, and hematoxylin counterstaining.<sup>21</sup>

*Western blot.* Western blots on digested lungs and BAL samples were done using standard techniques and immunoblotted with CD41 and α-tubulin Abs.

*ELISA.* 15-epi-LXA<sub>4</sub> and thromboxane B<sub>2</sub> concentrations in plasma and BAL were determined using ELISA kits from Neogen and GE Healthcare, respectively. We followed the manufacturer's instructions for lipid extraction.

*MPO activity.* Cell-free BAL and MPO standards (Sigma) were placed onto a 96-well plate and 200µl of o-Dianisidine solution and 10µl of 0.1% H<sub>2</sub>O<sub>2</sub> were added. The absorbance was read after 5 minutes at 405 nm, and expressed as units/ml of BAL supernatant using a standard curve.<sup>21</sup>

*Flow cytometry.* Whole blood was collected into acid citrate dextrose (ACD, Sigma). Samples in the presence of Fc $\gamma$ RII/III blocking Ab (2.4G2) were diluted with Tyrode's buffer supplemented with 10 units/mL of heparin (APP), 7 units/mL Apyrase (Sigma) and 0.5 $\mu$ M PGI<sub>2</sub> (Sigma). eFluor450-CD11b (eBioscience), PE-Ly6G (BD Bioscience), APC-CD41 (eBioscience), and isotope controls Abs were used to detect leukocyte and platelets antigens. Samples were examined with a LSRII/Fortessa flow cytometer (BD Bioscience). Neutrophils and monocytes were gated by their forward- and side-scatter characteristics, and by their Ly-6G<sup>+</sup>/CD11b<sup>+</sup> (neutrophil) or Ly-6G<sup>-</sup>/CD11b<sup>+</sup>/ Ly-6C<sup>+</sup> (monocyte) expression pattern. Platelet-neutrophil or platelet-monocyte aggregates were detected by CD41 Ab staining. In selected experiments, whole lungs were digested<sup>21</sup> followed by the same general protocol described above. All data were analyzed using FlowJo software (TreeStar).

*In vitro neutrophil-platelet aggregate experiments.* Whole blood was collected from C57BL/6 WT and *Fpr2/3<sup>-/-</sup>* mice into ACD, and used to isolate platelets.<sup>23</sup> Bone-marrow neutrophils were isolated, as previously described.<sup>24</sup> Neutrophils and platelets (1:100 ratio) from WT and *Fpr2/3<sup>-/-</sup>* mice were incubated for 1h at 37°C with LPS (5  $\mu$ g/mL) or with 15-epi-LXA<sub>4</sub> (300 nM), in presence of Tyrode's buffer. After this incubation, Tyrode's buffer with supplements (see above) was added. Flow cytometry was performed as described above.

*Lung intravital microscopy:* We used two-photon intravital microscopy<sup>25</sup> to record in real-time neutrophil and platelet interactions in LPS-induced ALI. To facilitate tracking of both platelets and neutrophils, *PF4-cre x Rosa26-LSL-tdTomato* mice (both from The Jackson Laboratories) were crossed with *LysM-eGFP* mice (obtained from E. Robey, University of California, Berkeley). To permit identification of the lung vasculature, Cascade-blue dextran (50 $\mu$ L of 25mg/mL; Life Technologies) was injected into the jugular vein. Mice were treated with LPS  $\pm$  ASA and compared with LPS untreated mice (PBS). We captured a 0.4 mm<sup>2</sup> x-y surface area at



40  $\mu\text{m}$  z-depth, capturing a complete image every 1 minute for 60 minutes. Images were analyzed using Imaris 7.6.1 software (Bitplane) to quantify mobile neutrophils (green) that co-labeled for platelets (red). The number of events was averaged per minute, and mean of the highest 20 values obtained were recorded as the number of neutrophil-platelet aggregates per  $0.1\text{ mm}^2$  of lung tissue at  $40\mu\text{m}$  z-depth.

*Statistics.* All *in vivo* and *in vitro* experiments were repeated a minimum of three independent times. Results are reported as both individual data points and mean  $\pm$  SD. To determine significance, 2-tailed Student's *t* test, ANOVA, and log-rank (Mantel-Cox) tests were used as appropriate (GraphPad PRISM version 5.0). *P* values of less than or equal to 0.05 were deemed to be significant.

*Study approval.* All experiments were approved by the Institutional Animal Care and Use Committee at the University of California, San Francisco.

## Results

*Intratracheal LPS induces platelet sequestration and activation within the lungs and aspirin attenuates these responses.* To determine the role of platelets in experimental ALI, we used an intratracheal LPS model of lung injury. Immunohistochemistry for the platelet specific marker CD41, 48 hours after intra-tracheal LPS instillation (5 µg/g), revealed intense sequestration of platelets within LPS-injured lungs, compared to PBS-instilled control animals (Figure 1A, B). We cannot rule out that areas of particularly intense CD41 staining at vessel junctions could indicate the presence of megakaryocytes. CD41 protein expression in total lung digest was analyzed by Western blot, and confirmed the sequestration of platelets within the lungs of LPS-challenged animals (Figure 1D). Thromboxane A<sub>2</sub> is a marker of platelet activation and is rapidly hydrolyzed to its inactive stable metabolite, thromboxane B<sub>2</sub> (TXB<sub>2</sub>). Plasma TXB<sub>2</sub> levels were increased after LPS challenge (Figure 1E). Pre-treatment with aspirin 24h and 2h before LPS challenge, prevented platelet accumulation in the lungs (Figure 1C, D) and decreased plasma TXB<sub>2</sub> levels (Figure 1E) compared to controls (DMSO).

*Aspirin decreases LPS-induced lung injury and alveolar platelet accumulation.* The total number of white blood cells (WBC) and neutrophils in bronchoalveolar lavage (BAL) was measured 48h after intratracheal PBS (control) or LPS instillation. MPO activity, an index of neutrophil activation, was also measured in cell-free BAL supernatant. The number of WBCs and neutrophils within BAL increased with LPS challenge, as did MPO activity (Figure 2A, B). BAL total protein was measured as an indicator of lung permeability, and increased after LPS instillation (Figure 2C). Mice treated with aspirin had significantly decreased BAL WBCs, neutrophils, MPO activity, and total protein, compared to vehicle controls (Figure 2A-C).

CD41 immunohistochemistry of lungs after LPS challenge revealed intense sequestration of platelets within the pulmonary capillaries, but also suggested the presence of intra-alveolar

platelets (Figure 1B inset). To further test for intra-alveolar platelet accumulation, we probed for CD41 protein via Western blot analysis on cells obtained from BAL. CD41 protein was increased in BAL from LPS-challenged mice, compared to PBS controls (Figure 2D). We also used flow cytometry and a CD41 mAb to determine the presence of platelets in the BAL, and did not detect intra-alveolar platelets in control animals, but a sharp increase was detected after LPS challenge (Figure 2E). We tested for pulmonary hemorrhage as a potential cause of intra-alveolar accumulation of platelets by measuring BAL hemoglobin and found a minimal increase in hemoglobin in the LPS-challenged animals (Supplemental Figure 1). We detected significantly increased TXB<sub>2</sub> concentrations in the cell-free-BAL supernatant of LPS-treated mice, compared to controls (Figure 2F). These results demonstrate that platelets accumulate and are activated in the alveolar spaces after LPS challenge. Aspirin treatment yielded a significant reduction in both alveolar LPS-induced CD41 expression (Figure 2D, E) and platelet activation (Figure 2F) and did not affect BAL hemoglobin measurements.

*Leukocyte-platelet aggregates are present in the blood and alveolar spaces after LPS-challenge and are reduced with aspirin treatment.* We investigated the LPS-induced formation of leukocyte-platelet aggregates (LPA) in whole blood and BAL using flow cytometry. The gating strategies for whole blood and BAL are illustrated in Supplemental Figure 2A-B. Briefly, antibodies against the leukocyte-specific surface markers CD11b, Ly6G or Ly6C, in addition to the platelet-specific marker CD41, were used to identify monocyte-platelet aggregates (MPA), defined as CD11b<sup>+</sup>/Ly6C<sup>+</sup>/Ly6G<sup>-</sup>/CD41<sup>+</sup>, and neutrophil-platelet aggregates (NPA), defined as CD11b<sup>+</sup>/Ly6G<sup>+</sup>/CD41<sup>+</sup>. Whole blood obtained by cardiac puncture revealed that LPS induced both NPA (Figure 2G) and MPA (Figure 2H) formation as early as 4h and 24h, respectively, after LPS instillation. Blood NPA and MPA levels progressively increased over time, reaching a maximum at 48h after LPS instillation (NPA = 62 ± 5.6%, MPA = 48.6 ± 3.4%). The percentage of NPA in the BAL of LPS-challenged mice also increased (Figure 2I), but we did not detect a

similar increase in BAL MPA (data not shown). Treatment with aspirin significantly decreased blood NPA and MPA formation, as well as BAL NPA formation, at 48h after LPS (Figure 2G-I).

*Dynamic formation of NPA within the lung microcirculation.* We used two-photon intravital lung microscopy<sup>25</sup> to determine the spatial and temporal formation of NPA within the pulmonary microcirculation at baseline and after LPS-induced ALI. To track platelets, *Rosa26-LSL-tdTomato* mice were crossed with *PF4-cre* mice,<sup>26</sup> and in the peripheral blood of these mice, >99% of the tdTomato<sup>+</sup> events were also CD41<sup>+</sup> (Supplemental Figure 3). These mice were crossed with *LysM-eGFP* mice to facilitate neutrophil identification. Cascade-blue dextran was injected i.v. to delineate the pulmonary microvasculature. In the triple transgenic mice without LPS exposure, we observed the dynamic formation of NPA within the lung microcirculation, with intravascular neutrophils interacting with platelets (example shown in Figure 3A and Supplemental Movie 1). We also observed NPA in PBS challenged mice (Figure 3B, E and Supplemental Movie 2), and in the first hours after LPS exposure there was a sharp increase in NPA formation (Figure 3C, F and Supplemental Movies 3 and 4), which was ameliorated with aspirin pre-treatment with NPA formation decreasing to basal levels (Figure 3D, G and Supplemental Movie 5). At later time points after LPS challenge (24 hours), we detected immobile NPA that were outside of the intravascular dextran signal (Figure 3H, I and Supplemental Movie 9), consistent with intra-alveolar NPA. Areas of neutrophil-platelet co-localization were analyzed by surface rendering (Supplemental Movies 6 – 8) and quantified in Figure 3J.

*Aspirin treatment increases plasma and BAL 15-epi-LXA<sub>4</sub> concentrations, and blockade of ALX reverses the protective effects of aspirin.* Plasma and BAL 15-epi-LXA<sub>4</sub> concentrations were measured by ELISA after PBS and LPS challenge ± aspirin treatment. LPS did not affect plasma or BAL 15-epi-LXA<sub>4</sub> levels, but aspirin treatment significantly increased both plasma and

BAL 15-epi-LXA<sub>4</sub> concentrations in LPS-challenged mice at 48h compared to PBS or vehicle controls (Figure 4A, B).

To determine whether elevated 15-epi-LXA<sub>4</sub> contributed to the decreased lung injury observed after aspirin treatment, we tested the ALX (Fpr2/3) antagonist, Boc2.<sup>27</sup> The administration of Boc2 30min before aspirin treatment, and repeated 24h after LPS instillation, reversed the protective effects of aspirin on lung injury (Figure 4C, D). Additionally, Boc2 partially reversed the decrease in whole blood NPA after aspirin treatment (Figure 4E). Boc2 treatment alone (in the absence of aspirin) had no effect on LPS-induced lung inflammation or injury (data not shown). Also, Boc2 did not change the plasma 15-epi-LXA<sub>4</sub> concentrations of aspirin treated mice (data not shown).

Next, we confirmed the results obtained with pharmacologic blockade of Fpr2/3 using the recently characterized *Fpr2/3*<sup>-/-</sup> mice.<sup>19</sup> Aspirin treatment of *Fpr2/3*<sup>-/-</sup> animals increased plasma concentrations of 15-epi-LXA<sub>4</sub> in the LPS model (Figure 4F). Compared to WT controls, *Fpr2/3*<sup>-/-</sup> mice had no significant differences in BAL WBCs, neutrophils or total protein after LPS challenge (data not shown), and we observed no significant decreases in BAL WBCs, neutrophils, or total protein after aspirin treatment (Figure 4G, H). The aspirin-induced decrease in blood and BAL NPA was also blocked in *Fpr2/3*<sup>-/-</sup> mice (Figure 4I, J).

*Effects of 15-epi-LXA<sub>4</sub> on LPS-induced lung injury.* To evaluate whether pharmacologic doses of 15-epi-LXA<sub>4</sub> can modulate LPS-induced lung injury, mice were treated with increasing doses of this lipid mediator. Intravenous administration of between 100-1500 ng of 15-epi-LXA<sub>4</sub> did not produce a significant change in BAL WBC counts or total protein after LPS challenge (Figure 5A, B). However, doses of 5 µg resulted in a significant decrease in BAL WBCs (Figure 5A) and

attenuated lung permeability (Figure 5B) and whole blood NPA formation (Figure 5C) in a similar manner to that observed with aspirin treatment.

To determine whether lipoxin receptor signaling on neutrophils and/or platelets mediated the protective effects observed after aspirin and 15-epi-LXA<sub>4</sub> treatment, we developed a co-culture system in which neutrophils and platelets, freshly isolated from WT or *Fpr2/3*<sup>-/-</sup> mice, were incubated with LPS (5 ug/mL) to induce NPA formation. We observed a basal formation of NPA after a 1h incubation of WT neutrophils and platelets (1:100), which increased after LPS treatment (Figure 5D). Pre-treatment with 15-epi-LXA<sub>4</sub> (300 nM) significantly decreased LPS-induced NPA formation (Figure 5D). We next tested combinations of WT and *Fpr2/3*<sup>-/-</sup> neutrophils and platelets, and observed that the presence of a functional lipoxin receptor was required on both neutrophils and platelets for 15-epi-LXA<sub>4</sub> to decrease NPA formation (Figure 5D).

*Lipoxins enhance inflammatory resolution in TRALI.* We have previously demonstrated that aspirin treatment decreases platelet sequestration, lung vascular permeability and edema formation, and increases survival in an experimental TRALI model.<sup>2,24</sup> Here, we investigated whether ATL was required for these effects. In a two-event model of TRALI,<sup>21</sup> we treated animals with aspirin ± Boc2, or vehicle (DMSO), 30min prior to the i.p. LPS priming, and again 2h prior to challenge with H2K<sup>d</sup> mAb. Both aspirin and aspirin + Boc2 treatment significantly increased 15-epi-LXA<sub>4</sub> concentrations in this model (Figure 6A). Lung edema formation, lung vascular permeability, hemoconcentration, and mortality were all reduced with aspirin treatment compared with vehicle-treated controls (Figure 6B-D, F). Extravascular lung water and EVPE values in LPS-primed controls are 0 and <10 µl, respectively.<sup>21</sup> Hematocrit values in LPS-primed controls are approximately 50%.<sup>21</sup> TRALI induced the formation of lung NPA, which was ameliorated by aspirin treatment (Figure 6E). Pretreatment with Boc2 significantly reversed the

protective effects of aspirin on lung injury, NPA formation, and mortality (Figure 6B-F). The administration of Boc2 alone did not affect TRALI severity (data not shown).

We subsequently tested the impact of treatment with 15-epi-LXA<sub>4</sub> or vehicle (i.v.) administered 30 min prior to H2K<sup>d</sup> mAb challenge on lung injury and NPA formation. 15-epi-LXA<sub>4</sub> treatment significantly decreased lung injury and formation of lung NPA, and eliminated mortality, compared to vehicle-treated animals (Figure 7A-E).

## Discussion

The main conclusions from this study are that (a) pulmonary platelet sequestration and activation, including intra-alveolar accumulation, are major features of ALI; (b) NPA are dynamically formed in the lung microcirculation, and sharply increase in the blood and alveolar compartments after injury; (c) ATL is required for aspirin-mediated protection in ALI, and ATL signals via Fpr2/3 on neutrophils and platelets to regulate the formation of NPA; and (d) in two experimental models of ALI, ATL treatment alone significantly reduces lung injury.

Platelets are capable of rapidly responding to inflammatory stimuli, releasing preformed mediators and even synthesizing immune mediators from mRNA transferred from megakaryocytes.<sup>28</sup> Platelets and platelet microparticles<sup>29</sup> can bind to leukocytes and the endothelium, influencing the function of these cells during inflammation. Indeed, using lung intravital imaging, we have now shown that the lung is a site of *de novo* NPA formation under basal conditions. The unique structural features of the lung microcirculation that influence neutrophil trafficking,<sup>30</sup> and the proposed role of the lung in thrombopoiesis,<sup>31</sup> may enhance these interactions. With inflammation, we found an early and rapid increase in dynamic NPA formation in the lung microcirculation and many of these aggregates were ultimately detected in the alveolar spaces. Previous reports of NPA formation in inflammation have shown that neutrophils bound to the endothelium secondarily capture platelets,<sup>32,33</sup> but our direct imaging of the lung microcirculation reveals that these aggregates are mobile, and are eventually capable of migrating into the alveolar spaces. It has recently been described that the PF4-cre mouse is potentially leaky in non-megakaryocyte lineages,<sup>44</sup> however, in the peripheral blood we found that >99% of the PF4-cre driven tdTomato<sup>+</sup> events were platelet specific (CD41<sup>+</sup>).

There is a current lack of understanding of the regulation of LPA formation. Here, we propose that anti-inflammatory lipoxins and ATL specifically, regulate this process. Lipoxins and ATL



exert potent anti-inflammatory and pro-resolution bio-actions. In particular, they block chemotaxis, adherence to microvasculature, inhibit NF- $\kappa$ B activation in leukocytes, and block the release of pro-inflammatory cytokines such as IL-6, IL-8 and TNF- $\alpha$ .<sup>16,34-37</sup> A potential limitation of our study is that microgram quantities of ATL were required to limit lung injury in our experiments, whereas others have reported that nanogram amounts of pro-resolving lipid mediators are sufficient.<sup>14</sup> However, we treated with ATL at just two time points in our 48 hour LPS model, and more frequent dosing of ATL, which has a half-life of a few minutes, may have allowed for lower overall amounts. Also, we have not ruled out the contribution of other aspirin-triggered lipid mediators, such as resolvins, but results from our pharmacologic experiments support a major role of ATL in mediating the protective effects of aspirin.

Others have shown protective effects of lipoxins in acute inflammation. LXA<sub>4</sub> and ATL were protective in an acid-induced ALI model through the modulation of neutrophil apoptosis.<sup>38</sup> ATL was also protective in mice challenged with LPS through a heme-oxygenase1-dependent mechanism.<sup>39</sup> Endogenous LXA<sub>4</sub> modulates ischemia-reperfusion injury in the gut, and *Fpr2/3*<sup>-/-</sup> mice have more inflammation and injury compared to wild-type animals.<sup>40</sup> In this same study, NPA formation was investigated using chimeric experiments to test whether platelet or neutrophil *Fpr2/3* was more important in aggregate development, and neutrophil *Fpr2/3* was determined to be critical.<sup>40</sup> However, in our neutrophil-platelet aggregation assay, we found that neither neutrophil nor platelet *Fpr2/3* is required for the formation of aggregates in response to LPS, but *Fpr2/3* on both cells is required for lipoxin-mediated inhibition of NPA formation.

For the first time, we have shown the presence of platelets in the alveolar spaces in ALI. Others have shown in experimental allergic inflammation, that platelets migrate out of vessels and localize beneath airways<sup>41</sup> or into the synovial spaces in rheumatoid arthritis.<sup>8</sup> Platelets have also been observed by electron microscopy extravasating out of the lung vasculature in patients

with ARDS.<sup>42</sup> In our studies, many of the platelets were bound to neutrophils, and the dynamics of NPA migration into the alveolar spaces after LPS challenge seems to occur hours after the very early formation of NPA in the lung vessels. More work is needed to determine the mechanisms of platelet transmigration in the lung, including the chemotactic stimuli for platelets, and whether transmigration is facilitated by tethering to neutrophils or occurs independently. Notably, our assays could have detected intact platelets or platelet microparticles in the alveolar spaces, and the specific role of platelet microparticles in lung inflammation should be addressed in future studies.

We previously reported that aspirin is protective in a two-event model of TRALI by decreasing platelet sequestration and activation and NET formation.<sup>2,21</sup> Here, we found that ATL and ALX were required for the protective effects of aspirin in TRALI. Indeed, treatment of mice with ATL alone yielded significant protection from TRALI. These results have important clinical implications for patients with ARDS including transfused, critically ill patients. Platelet depletion is impractical, as is aspirin therapy in many critically ill patients, but treatment with ATL, or perhaps other anti-inflammatory and pro-resolving mediators,<sup>43</sup> is an attractive approach. Further, the measurement of LPA in patients with ARDS and other critical illnesses could be a useful biomarker of inflammation, and could be measured serially to assess therapeutic responses to treatment with pro-resolving lipid mediators.

In summary, we have shown in two experimental models of ALI that aspirin-triggered lipoxin is a powerful inhibitor of neutrophil- and platelet-mediated lung inflammation and injury. Ironically, surface contact between neutrophil and platelets yields transcellular production of lipoxins and also leukotrienes,<sup>45</sup> and our results indicate that signaling through ALX on neutrophils and platelets in turn regulates their tethering during inflammation. Aspirin is likely exerting its therapeutic benefit through a combination of inhibiting the production of pro-inflammatory

eicosanoids, and triggering the formation of pro-resolving lipid mediators.<sup>46</sup> Even low-dose dose aspirin administered to healthy volunteers is capable of producing bioactive levels of ATL,<sup>47</sup> and a clinical trial (ClinicalTrials.gov Identifier: NCT01504867) is underway randomizing patients at risk for ARDS to aspirin or placebo. ATL and potentially other pro-resolving lipid mediators are attractive therapeutic options for patients with ARDS—a life-threatening syndrome that lacks any effective pharmacotherapies.

## **Acknowledgments**

We thank K. Corbin and H. Pinkard at the UCSF Biological Imaging Development Center for assistance with the intravital lung microscopy experiments. We also thank Rod Flower at the William Harvey Research Institute, St. Barts and the London School of Medicine and Dentistry for providing the *Fpr2/3<sup>-/-</sup>* mice. This work was supported by US National Institutes of Health grants R01 HL107386 (M.R.L.) and P01 HL024136 (M.F.K).

## **Authorship Contributions**

G.O-M. designed and performed the experiments, analyzed the results, and prepared the manuscript.

B.M. performed experiments and edited the manuscript.

A.B. assisted with the intravital lung imaging experiments and edited the manuscript.

M.H. assisted with the intravital lung imaging experiments and edited the manuscript.

M.F.K. assisted with the intravital lung imaging experiments and edited the manuscript.

M.R.L. designed and assisted with the experiments, analyzed the results, provided funding, and prepared and edited the manuscript.

## **Disclosure of Conflicts of Interest**

None of the authors have competing financial interests.

## References

1. Matthay MA, Ware LB, Zimmerman GA. The acute respiratory distress syndrome. *J Clin Invest.* 2012; 122(8):2731-2740.
2. Caudrillier A, Kessenbrock K, Gilliss BM, et al. Platelets induce neutrophil extracellular traps in transfusion-related acute lung injury. *J Clin Invest.* 2012; 122(7):2661-2671.
3. Vieira-de-Abreu A, Campbell RA, Weyrich AS, et al. Platelets: versatile effector cells in hemostasis, inflammation, and the immune continuum. *Semin Immunopathol.* 2012; 34(1):5-30.
4. Zarbock A, Singbartl K, Ley K. Complete reversal of acid-induced acute lung injury by blocking of platelet-neutrophil aggregation. *J Clin Invest.* 2006; 116(12):3211-3219.
5. Cerletti C, Tamburrelli C, Izzi B, et al. Platelet-leukocyte interactions in thrombosis. *Thromb Res.* 2012; 129(3):263-266.
6. Gawaz M, Fateh-Moghadam S, Pilz G, et al. Platelet activation and interaction with leucocytes in patients with sepsis or multiple organ failure. *Eur J Clin Invest.* 1995; 25(11):843-851.
7. Irving PM, Macey MG, Shah U, et al. Formation of platelet-leukocyte aggregates in inflammatory bowel disease. *Inflamm Bowel Dis.* 2004; 10(4):361-372.
8. Boilard E, Nigrovic PA, Larabee K, et al. Platelets amplify inflammation in arthritis via collagen-dependent microparticle production. *Science.* 2010; 327(5965):580-583.
9. O'Sullivan BP, Linden MD, Frelinger AL, 3rd, et al. Platelet activation in cystic fibrosis. *Blood.* 2005; 105(12):4635-4641.
10. Furman MI, Benoit SE, Barnard MR, et al. Increased platelet reactivity and circulating monocyte-platelet aggregates in patients with stable coronary artery disease. *J Am Coll Cardiol.* 1998; 31(2):352-358.

11. Claria J, Serhan CN. Aspirin triggers previously undescribed bioactive eicosanoids by human endothelial cell-leukocyte interactions. *Proc Natl Acad Sci U S A*. 1995; 92(21):9475-9479.
12. Serhan CN. Lipoxins and aspirin-triggered 15-epi-lipoxins are the first lipid mediators of endogenous anti-inflammation and resolution. *Prostaglandins Leukot Essent Fatty Acids*. 2005; 73(3-4):141-162.
13. Levy BD, Clish CB, Schmidt B, et al. Lipid mediator class switching during acute inflammation: signals in resolution. *Nat Immunol*. 2001; 2(7):612-619.
14. Serhan CN. Resolution phase of inflammation: novel endogenous anti-inflammatory and proresolving lipid mediators and pathways. *Annu Rev Immunol*. 2007; 25:101-137.
15. Serhan CN, Sheppard KA. Lipoxin formation during human neutrophil-platelet interactions. Evidence for the transformation of leukotriene A4 by platelet 12-lipoxygenase in vitro. *J Clin Invest*. 1990; 85(3):772-780.
16. Serhan CN, Maddox JF, Petasis NA, et al. Design of lipoxin A4 stable analogs that block transmigration and adhesion of human neutrophils. *Biochemistry*. 1995; 34(44):14609-14615.
17. Chiang N, Serhan CN, Dahlen SE, et al. The lipoxin receptor ALX: potent ligand-specific and stereoselective actions in vivo. *Pharmacol Rev*. 2006; 58(3):463-487.
18. Czapiga M, Gao JL, Kirk A, et al. Human platelets exhibit chemotaxis using functional N-formyl peptide receptors. *Exp Hematol*. 2005; 33(1):73-84.
19. Dufton N, Hannon R, Brancaleone V, et al. Anti-inflammatory role of the murine formyl-peptide receptor 2: ligand-specific effects on leukocyte responses and experimental inflammation. *J Immunol*. 2010; 184(5):2611-2619.
20. Cooray SN, Gobbetti T, Montero-Melendez T, et al. Ligand-specific conformational change of the G-protein-coupled receptor ALX/FPR2 determines proresolving functional responses. *Proc Natl Acad Sci U S A*. 2013; 110(45):18232-18237.

21. Looney MR, Nguyen JX, Hu Y, et al. Platelet depletion and aspirin treatment protect mice in a two-event model of transfusion-related acute lung injury. *J Clin Invest.* 2009; 119(11):3450-3461.
22. Bozza FA, Shah AM, Weyrich AS, et al. Amicus or adversary: platelets in lung biology, acute injury, and inflammation. *Am J Respir Cell Mol Biol.* 2009; 40(2):123-134.
23. Krueger LA, Barnard MR, Frelinger AL, 3rd, et al. Immunophenotypic analysis of platelets. *Curr Protoc Cytom.* 2002; Chapter 6:Unit 6 10.
24. Looney MR, Su X, Van Ziffle JA, et al. Neutrophils and their Fc gamma receptors are essential in a mouse model of transfusion-related acute lung injury. *J Clin Invest.* 2006; 116(6):1615-1623.
25. Looney MR, Thornton EE, Sen D, et al. Stabilized imaging of immune surveillance in the mouse lung. *Nat Methods.* 2011; 8(1):91-96.
26. Tiedt R, Schomber T, Hao-Shen H, et al. Pf4-Cre transgenic mice allow the generation of lineage-restricted gene knockouts for studying megakaryocyte and platelet function in vivo. *Blood.* 2007; 109(4):1503-1506.
27. von der Weid PY, Hollenberg MD, Fiorucci S, et al. Aspirin-triggered, cyclooxygenase-2-dependent lipoxin synthesis modulates vascular tone. *Circulation.* 2004; 110(10):1320-1325.
28. Rondina MT, Weyrich AS, Zimmerman GA. Platelets as cellular effectors of inflammation in vascular diseases. *Circ Res.* 2013; 112(11):1506-1519.
29. Italiano JE, Jr., Mairuhu AT, Flaumenhaft R. Clinical relevance of microparticles from platelets and megakaryocytes. *Curr Opin Hematol.* 2010; 17(6):578-584.
30. Burns AR, Smith CW, Walker DC. Unique structural features that influence neutrophil emigration into the lung. *Physiol Rev.* 2003; 83(2):309-336.
31. Weyrich AS, Zimmerman GA. Platelets in lung biology. *Annu Rev Physiol.* 2013; 75:569-591.

32. Hidalgo A, Chang J, Jang JE, et al. Heterotypic interactions enabled by polarized neutrophil microdomains mediate thromboinflammatory injury. *Nat Med.* 2009; 15(4):384-391.
33. Jenne CN, Wong CH, Petri B, et al. The use of spinning-disk confocal microscopy for the intravital analysis of platelet dynamics in response to systemic and local inflammation. *PLoS One.* 2011; 6(9):e25109.
34. Colgan SP, Serhan CN, Parkos CA, et al. Lipoxin A4 modulates transmigration of human neutrophils across intestinal epithelial monolayers. *J Clin Invest.* 1993; 92(1):75-82.
35. Scalia R, Gefen J, Petasis NA, et al. Lipoxin A4 stable analogs inhibit leukocyte rolling and adherence in the rat mesenteric microvasculature: role of P-selectin. *Proc Natl Acad Sci U S A.* 1997; 94(18):9967-9972.
36. Jozsef L, Zouki C, Petasis NA, et al. Lipoxin A4 and aspirin-triggered 15-epi-lipoxin A4 inhibit peroxynitrite formation, NF-kappa B and AP-1 activation, and IL-8 gene expression in human leukocytes. *Proc Natl Acad Sci U S A.* 2002; 99(20):13266-13271.
37. Sodin-Semrl S, Taddeo B, Tseng D, et al. Lipoxin A4 inhibits IL-1 beta-induced IL-6, IL-8, and matrix metalloproteinase-3 production in human synovial fibroblasts and enhances synthesis of tissue inhibitors of metalloproteinases. *J Immunol.* 2000; 164(5):2660-2666.
38. El Kebir D, Jozsef L, Pan W, et al. 15-epi-lipoxin A4 inhibits myeloperoxidase signaling and enhances resolution of acute lung injury. *Am J Respir Crit Care Med.* 2009; 180(4):311-319.
39. Jin SW, Zhang L, Lian QQ, et al. Posttreatment with aspirin-triggered lipoxin A4 analog attenuates lipopolysaccharide-induced acute lung injury in mice: the role of heme oxygenase-1. *Anesth Analg.* 2007; 104(2):369-377.
40. Brancaleone V, Gobbetti T, Cenac N, et al. A vasculo-protective circuit centered on lipoxin A4 and aspirin-triggered 15-epi-lipoxin A4 operative in murine microcirculation. *Blood.* 2013; 122(4):608-617.
41. Pitchford SC, Momi S, Baglioni S, et al. Allergen induces the migration of platelets to lung tissue in allergic asthma. *Am J Respir Crit Care Med.* 2008; 177(6):604-612.



42. Bachofen M, Weibel ER. Structural alterations of lung parenchyma in the adult respiratory distress syndrome. *Clin Chest Med.* 1982; 3(1):35-56.
43. Levy BD, Serhan CN. Resolution of acute inflammation in the lung. *Annu Rev Physiol.* 2014; 76:467-492.
44. Calaminus SD, Guitart AV, Sinclair A, et al. Lineage tracing of Pf4-Cre marks hematopoietic stem cells and their progeny. *PLoS One.* 2012; 7(12):e51361.
45. Fiore S, Serhan CN. Formation of lipoxins and leukotrienes during receptor-mediated interactions of human platelets and recombinant human granulocyte/macrophage colony-stimulating factor-primed neutrophils. *J Exp Med.* 1990; 172(5):1451-1457.
46. Eickmeier O, Seki H, Haworth O, et al. Aspirin-triggered resolvin D1 reduces mucosal inflammation and promotes resolution in a murine model of acute lung injury. *Mucosal Immunol.* 2013; 6(2):256-266.
47. Chiang N, Bermudez EA, Ridker PM, et al. Aspirin triggers antiinflammatory 15-epi-lipoxin A4 and inhibits thromboxane in a randomized human trial. *Proc Natl Acad Sci U S A.* 2004; 101(42):15178-15183.

## Figure Legends

**Figure 1. Platelets are sequestered and activated in the lungs after intratracheal LPS and aspirin attenuates these responses.** CD41 immunostaining of lung tissue sections from (A) control mice (PBS), (B) LPS-challenged mice treated with vehicle (DMSO), or (C) LPS-challenged mice treated with ASA. Images are representative of at least 3 animals per group. Original magnification x60 and zoom of outlined areas. Scale bar = 20  $\mu\text{m}$ . Arrows indicate CD41<sup>+</sup> events in the intra-alveolar spaces. (D) CD41 protein expression and densitometric analysis from lung digestion samples in PBS controls and LPS-challenged mice  $\pm$  ASA.  $^{\dagger\dagger\dagger}P < 0.001$  vs. PBS;  $^{****}P < 0.0001$  vs. vehicle. (E) Platelet activation was assessed by measuring thromboxane B<sub>2</sub> (TXB<sub>2</sub>) plasma levels in PBS and LPS-challenged mice  $\pm$  ASA.  $^{\dagger}P < 0.05$  vs. PBS;  $^{****}P < 0.0001$  vs. vehicle. Data are mean  $\pm$  SD of four to nine animals per group.

**Figure 2. LPS produces ALI, increased blood NPA and MPA, and the intra-alveolar accumulation of platelets and NPA; aspirin attenuates these responses.** (A) BAL white blood cell (WBC) and neutrophil counts, (B) MPO activity in the cell-free BAL, and (C) lung protein permeability in PBS and LPS-challenged mice + ASA vs. vehicle (DMSO) controls.  $^{**}P < 0.01$ ,  $^{****}P < 0.0001$  vs. vehicle groups and  $^{\dagger\dagger\dagger}P < 0.001$ ,  $^{\dagger\dagger\dagger\dagger}P < 0.0001$  vs PBS. (D) Densitometric analysis and representative membrane of blots against CD41 protein performed on BAL cells in PBS controls and LPS + ASA vs. vehicle group.  $^{\dagger\dagger\dagger\dagger}P < 0.0001$ ,  $^{****}P < 0.0001$  vs. PBS and vehicle, respectively. (E) CD41 (and isotype control) flow cytometry analysis in BAL cells from mice with no LPS and LPS  $\pm$  ASA with representative histogram plots and MFI (representative of 5 independent experiments). (F) TXB<sub>2</sub> levels in cell-free BAL from control and LPS + ASA vs. vehicle groups.  $^{\dagger}P < 0.05$  vs. PBS;  $^{**}P < 0.05$  vs. vehicle. (G) % neutrophil-platelet

aggregates (NPA) and (H) % monocyte-platelet aggregates (MPA) in whole blood at different time points after LPS instillation and effects of ASA treatment at indicated times.  $\dagger\dagger P < 0.01$ ,  $\dagger\dagger\dagger P < 0.001$ ,  $\dagger\dagger\dagger\dagger P < 0.0001$  vs. PBS;  $**P < 0.01$ ,  $****P < 0.0001$  vs. LPS-challenged mice at 48h. (I) %NPA in BAL after LPS + ASA vs. vehicle (at 48h).  $*P < 0.05$  vs. vehicle. Data are mean  $\pm$  SD of five to ten animals per group.

**Figure 3. Lung intravital microscopy reveals the dynamic formation of NPA in the lungs under homeostatic and injury conditions.** (A) Screenshots of a *PF4-cre x Rosa26-LSL-tdTomato x LysM-eGFP* mouse under basal conditions showing the dynamic formation of NPA in the lung microcirculation (see also Supplementary Movie 1). (B-G) Screenshots with zoom insets of *PF4-cre x Rosa26-LSL-tdTomato x LysM-eGFP* mice challenged i.t. with PBS (B, E), LPS + vehicle (C, F), or LPS + ASA (D, G), and imaged directly after challenge (see also Supplementary Movies 2-8). White arrows indicate NPA. (H-I) Screenshots from *PF4-cre x Rosa26-LSL-tdTomato x LysM-eGFP* mice challenged with i.t. LPS and imaged 24 hours after instillation. White arrows indicate intra-alveolar NPA (see also Supplementary Movie 9). (J) Quantification of NPA in B-D.  $\dagger P < 0.05$  vs. PBS;  $*P < 0.05$  vs. vehicle. Data are mean  $\pm$  SD of three to seven animals per group.

**Figure 4. Blocking the lipoxin receptor reverses aspirin protective effects.** (A) 15-epi-LXA<sub>4</sub> levels in plasma and (B) cell-free BAL of wild-type (WT) mice challenged with PBS or LPS + ASA vs. vehicle (DMSO).  $****P < 0.0001$  vs. PBS and vehicle groups. (C) BAL WBC and neutrophil counts, (D) BAL cell-free total protein and (E) whole blood NPA in WT mice challenged with LPS + ASA vs. vehicle or ASA + Boc2.  $***P < 0.001$  and  $****P < 0.0001$  vs. vehicle groups, respectively.  $\dagger P < 0.05$  and  $\dagger\dagger P < 0.01$  vs. ASA + Boc2 groups, respectively. (F)

15-epi-LXA<sub>4</sub> levels in plasma of *Fpr2/3*<sup>-/-</sup> mice challenged with PBS or LPS + ASA vs. vehicle (DMSO). \*\*\*\**P* <0.0001 vs. PBS and vehicle groups. (G) BAL WBC and neutrophil counts, (H) BAL cell-free total protein, (I) NPA in whole blood, and (J) NPA in BAL cells of *Fpr2/3*<sup>-/-</sup> mice challenged with PBS or LPS + ASA vs. vehicle. \**P* <0.05, \*\**P* <0.01 vs. PBS. NS = not significant. Data are mean ± SD of five to seven animals per group.

**Figure 5. Treatment with 15-epi-LXA<sub>4</sub> decreases LPS-induced ALI and regulates NPA formation in vivo and in vitro.** (A,B) ASA or 15-epi-LXA<sub>4</sub> treatment (at doses indicated) 2h before and 24h after LPS challenge and (A) leukocyte recruitment into the air spaces and (B) BAL total protein. \*\*\**P* <0.001, \*\*\*\**P* <0.0001 vs. vehicle groups. (C) Whole blood NPA in mice challenged with LPS ± ASA or 15-epi-LXA<sub>4</sub> (5 µg). \*\**P* <0.01, \*\*\*\**P* <0.0001 vs. vehicle. (D) Neutrophils and platelets isolated from WT or *Fpr2/3*<sup>-/-</sup> mice were incubated with LPS (5 µg/ml) for 1h to produce NPAs in vitro. LPS induced NPA in all combinations of WT and *Fpr2/3*<sup>-/-</sup> neutrophils/platelets. \**P* <0.05 vs. basal (no LPS). Pre-treatment with 15-epi-LXA<sub>4</sub> reduced NPA formation after LPS in only the WT neutrophil/WT platelet group. †*P* <0.05. Data are mean ± SD of five animals per group (A-C) or mean ± SD of at least three independent experiments (D).

**Figure 6. Boc2 blocks the protective effects of aspirin in a two-event model of TRALI.** BALB/c WT mice were treated with LPS (0.1 mg/kg, i.p.) and 24h later were challenged with H2K<sup>d</sup> mAb (1.0 mg/kg, i.v.) and sacrificed 2h later. (A) Plasma 15-epi-LXA<sub>4</sub> levels in mice with TRALI + vehicle, ASA, or ASA + Boc2. \*\*\*\**P* <0.0001 vs. vehicle group. (B) Extravascular lung water, (C) extravascular plasma equivalents (EVPE), and (D) hematocrit in mice with TRALI + vehicle, ASA, or ASA + Boc2. \**P* <0.05, \*\**P* <0.01, and \*\*\*\**P* <0.0001 vs. vehicle groups. †*P*

<0.05 and  $^{+++}P < 0.0001$  vs. ASA + Boc2 groups. (E) NPA in the lung digestion in mice with TRALI + vehicle, ASA, or ASA + Boc2.  $^*P < 0.05$  and  $^{***}P < 0.001$  vs. vehicle group.  $^{\dagger}P < 0.05$  vs. ASA + Boc2 group. (F) Survival curves in mice with TRALI + vehicle, ASA, or ASA + Boc2.  $^{**}P < 0.01$  vs. vehicle group and  $^{\dagger}P < 0.05$  vs. ASA + Boc2 group. Data are mean  $\pm$  SD of five to ten animals per group.

**Figure 7. 15-epi-LXA<sub>4</sub> protects mice from TRALI.** (A) Extravascular lung water, (B) extravascular plasma equivalent (EVPE), (C) NPA in the lung digestion and (D) hematocrit in WT mice with TRALI + 15-epi-LXA<sub>4</sub> vs. vehicle.  $^*P < 0.05$ ,  $^{**}P < 0.01$ , and  $^{***}P < 0.001$  vs. vehicle groups. (E) Survival curves in mice with TRALI + 15-epi-LXA<sub>4</sub> vs. vehicle.  $^*P < 0.05$  vs. vehicle group. Data are mean  $\pm$  SD of ten animals per group.

Figure 1







Figure 2




Figure 3






Figure 4





Figure 5




Figure 6




Figure 7

